



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

JOURNAL

OF THE

New York Entomological Society.

Vol. XI.

SEPTEMBER, 1903.

No. 3.

THE SKEWNESS OF THE THORAX IN THE ODONATA.

BY JAMES G. NEEDHAM AND MAUDE H. ANTHONY.

(PLATE VIII.)

Any one looking carefully at a dragonfly sees that the legs are attached far forward and the wings far back upon the thorax, and that the side plates of the latter are decidedly aslant. This arrangement of parts is an adaptation to perching on the sides of vertical stems without much alteration of the position maintained in flight. It makes for celerity in stopping and starting again. The legs are thrown forward where they readily reach and grasp the vertical stem, and the wings are shifted backward and tilted so that their cutting edges are directed obliquely upward, in which position a simple sculling action lifts the body instantly from its support.

In the jumping Orthoptera exactly the reverse inclination of the lateral thoracic sclerites has taken place: the legs have been shifted backward — especially the large hind ones used in jumping — and the side pieces are aslant with the opposite inclination. Doubtless these lateral sclerites (episternum and epimeron) were primitively placed at right angles to the axis of the body, so that the sutures between them were vertical, as they still are when first developed in dragonfly and grasshopper alike.

Among the orders the Odonata are extremely isolated, and, in their own way, undoubtedly highly specialized. As marks of their isolation the accessory genitalia of the males developed in an isolated position on the ventral side of the second abdominal segment, the

type of venation, and the remarkable structure of the labium — especially, of the nymphal labium — have been frequently noted. But this skewness of the thorax, hitherto almost unstudied, is the external evidence of the most profound alterations of the whole bodily organization. As for the skeleton, the legs have moved forward and the wings backward, greatly increasing the areas between the sternum of the metathorax and the abdomen, and between the tergum of the mesothorax and the prothorax respectively, and these areas have been overgrown by neighboring lateral sclerites (mesepisternum in front and metepimeron behind). The unusual proportions and the new (dorsal and ventral) positions thus attained by these sclerites were long a puzzle to many eminent entomologists. The question of their homologies was finally set at rest by a study of the segmental muscles and sutures made by Dr. Calvert for his well-known catalogue, published in 1893. He showed that the muscles have retained fully their segmental arrangement, the wing muscles becoming enormously enlarged and taking on the general inclination of the thorax. The mid-lateral suture is completely and the others are almost obliterated.

This fusion of sclerites is doubtless an accompaniment of the increasing power of the wing muscles. The skeleton is further strengthened by the development of a unique system of carinæ, the strongest of which is the mid-dorsal thoracic carina, formed at the junction of the mesepisterna along the dorsal line, forking above and ending in an antealar crest, ending below in a transverse collar-like ridge abutting against the prothorax. There are also carinæ along the upper ends of the lateral sclerites about the wing bases, and others trussing the floor of the metathorax between the bases of the hind legs and the abdomen. Doubtless these all contribute to the strength of the thoracic skeleton, and enable it to withstand the pull of the enormously large and powerful wing muscles. If in a dragonfly that has newly emerged from the nymphal skin and that has not yet had time for the hardening of the skeleton, the muscles be stimulated artificially to contract (as by putting in alcohol or cyanide bottle) they draw the thorax into a crumpled and contorted condition. Doubtless a careful study of this system of carinæ, and of the external topography of the thoracic skeleton in general would yield good results: but it is a less ambitious undertaking that this paper records.

Impressed by the differences in degree of skewness in the thorax of a number of dragonflies that were lying before him one day, the

senior author bethought himself that this skewness might be measured, and devised as an instrument for the purpose the goniometer shown in Pl. VIII, Fig. 1. This was constructed with little trouble out of a discarded box top, about 100 mm. square, a small brass protractor scale, a bicycle spoke, and a piece of brass about 25 mm. square. The brass was first drilled through the center and reamed out so that the head of the bicycle spoke would fit it neatly and rotate in it smoothly. Then the corners of the brass were drilled to receive screws. Then, with the spoke in place, its head flush with the surface of the brass, the latter was screwed fast to the under side of the wooden base, nearer the hinge edge of the cover, from which the side strip that was underneath had been removed. Thus the spoke was securely held by its head while free to rotate in the brass. Then the spoke was bent twice at right angles in an elongate **U** with unequal arms, the first bend perpendicular to, the second parallel to the surface of the board, the two arms being strictly parallel and far enough apart to allow the placing of the body of the largest dragonfly between the upper arm and the wooden base (Pl. VIII, Fig. 1). Then the protractor scale was glued to the wooden base in such position that its center of curvature was exactly over the center of the pivot below the base. The longer upper arm of the **U** then crossed the center in any position of rotation, and its end crossing the scale served as an indicator.

To use this goniometer a dragonfly with wings folded back to back was laid on a broad glass slip (this merely for convenience in moving the specimen) and brought to rest with its predetermined base line of angle measurement coinciding with the base line of the protractor scale. Then the index arm above was moved parallel with the suture forming the other limb of the angle to be measured. Then the angle was read by sighting along the edge of the indicator, keeping the exact center and the degree to be read in alignment. Thus the three successive operations — the placing of the specimen, the adjustment of the indicator and the reading of the scale — were done independently and in the order stated. This made for accuracy, but there were both mechanical and anatomical reasons why great accuracy was unattainable.

1. *Mechanical*. — The base line was too short. It was impossible to go beyond the confines of the combined meso- and metathorax and have fixed points, owing to the flexibility of the articulations with prothorax and with abdomen. To settle upon two points that should

determine a longitudinal line in comparison with which the angle of inclination of the sclerites should be measured was not easy. After canvassing the external topography of the thorax carefully, we settled upon the pleural articulation of the middle coxa for the anterior point (Pl. VIII, Fig. 2, *b*) and the infero-lateral articulation of the thorax and abdomen for the posterior, the two determining the base line *bd* of Fig. 2. By comparing Fig. 1* it will be seen that the two points are so close together as to occasion difficulty in bringing them into exact coincidence with the base line of the scale.

2. *Anatomical*. — The anatomical sources of error were several. (*a*) The articulations used to determine the base line are something more than points in breadth. (*b*) They are sometimes obscured by hairs. (*c*) The sutures with which the indicator must be made parallel are sometimes sinuous, and their general direction has to be estimated.

At first the skewness of the three lateral sutures and of the dorsal carina were measured, but as the differences discovered were rather less than the rather wide limits of probable error, only the first lateral (humeral) suture was measured to the end, and that and the tilt of the wing bases in the opposite direction are reported upon below.

The diagrammatized photograph (Fig. 2) shows these angles: *abc* is the angle made by the humeral suture *cb* with the perpendicular *ab* to the base line *db*, assumed to be parallel with the axis of the body. This angle measured upon the arc *x* represents the degree of skewness or inclination of that suture.

cdb is the angle made by a line *cd* drawn through the wing bases with the base line *bd*, and is measured upon the arc *z*. The wing bases are assumed to have rested primitively upon the line *ec* parallel to *db*. The specimen shown in Fig. 1 is nearly in position for measuring this angle.

The actual measurements were all made by the junior author upon miscellaneous papered specimens. Each specimen selected was measured first upon one side and then upon the other, and after intervening measurement of other species, was measured again, and then the average of all the measurements was taken. But one specimen was used for each species and the sex was disregarded. Some of the first meas-

* The specimen is off the base line in Fig. 1, having slipped out of place just before this photograph was taken.

urements made showed considerable discrepancies and were discarded altogether; but, with practice, they came into much closer agreement. It is not claimed, however, that the figures herein given are to be relied upon absolutely within the limits of two or three degrees.

Suborder **Anisoptera.**

Fam. **ÆSCHNIDÆ.**

ÆSCHNINÆ.

	<i>x</i>	<i>z</i>		<i>x</i>	<i>z</i>
<i>Boyeria irene</i>	21	19*	<i>Staurophlebia magnifica</i> ...	36	25
<i>Anax junius</i>	27	21	<i>Epiæschna heros</i>	37	23
<i>Æschna californica</i> ..	34	22	<i>Gomphæschna furcillata</i> ...	37	24
<i>Coryphæschna ingens</i>	34	25	<i>Gynacantha trifida</i>	39	27
<i>Æschnophlebia anisoptera</i>	35	34†	<i>Basiæschna janata</i>	42	27
<i>Æschna verticalis</i>	36	23	<i>Planæschna multipunctata</i>	42	28

GOMPHINÆ.

<i>Dromogomphus spinosus</i>	31	25	<i>Lanthus albistylus</i>	42	28
<i>Gomphus simillimus</i> ..	32	22	<i>Herpetogomphus designatus</i>	41	28
<i>Gomphus villosipes</i>	33	26	<i>Ophiogomphus serpentinus</i> .	44	27
<i>Gomphoides stigmatus</i>	36	26	<i>Progomphus obscurus</i>	44	27
<i>Gomphidia confluens</i>	37	26	<i>Onychogomphus uncatus</i> ...	47	27
<i>Aphylla producta</i>	39	27	<i>Epigomphus paludosus</i>	47	29
<i>Hagenius brevistylus</i>	40	22*	<i>Hemigomphus ochraceus</i>	50	29†
<i>Gomphus dilatatus</i> ..	41	22			

PETALURINÆ.

<i>Tachopteryx thoreyi</i>	40	28
----------------------------------	----	----

CORDULEGASTERINÆ.

<i>Cordulegaster diastatops</i>	37	35
---------------------------------------	----	----

Fam. **LIBELLULIDÆ.**

MACROMIINÆ.

<i>Didymops transversa</i>	29	20	<i>Epophthalmia elegans</i>	33	18*
<i>Macromia magnifica</i>	32	22	<i>Synthemis brevistyla</i>	39	33†

* Minima: second column.

† Maxima: second column.

CORDULINÆ.

<i>Helocordulia uhleri</i>	30	25	<i>Tetragoneuria canis</i>	36	29†
<i>Epicordulia princeps</i>	33	24	<i>Cordulia shurtleffi</i>	37	24
<i>Hemicordulia tau</i>	34	23*	<i>Somatochlora elongata</i>	38	24

LIBELLULINÆ.

<i>Perithemis domitia</i>	28	21	<i>Mesothemis simplicicollis</i>	43	34
<i>Acisoma panorpoides</i>	28	24	<i>Trithemis minuscula</i>	43	38*
<i>Rhyothemis splendida</i>	33	25	<i>Brachythemis contaminata</i> ..	44	26
<i>Onychothemis abnormis</i>	31	22	<i>Tramea carolina</i>	44	27
<i>Brechmorhoga mendax</i>	34	24	<i>Pachydiplax longipennis</i>	44	27
<i>Lepthemis vesiculosa</i>	34	27	<i>Crocothemis erythræa</i>	44	30
<i>Belonia herculea</i>	35	19*	<i>Anatya anomala</i>	45	29
<i>Diastatops tincta</i>	36	27	<i>Neurothemis equestris</i>	45	31
<i>Zygonyx iris</i>	36	28	<i>Micrathyria didyma</i>	47	27
<i>Dythemis velox</i>	37	29	<i>Potamothemis americana</i> ..	46	29
<i>Orthetrum albistylum</i>	38	25	<i>Sympetrum rubicundulum</i> ..	46	31
<i>Miathyria marcella</i>	39	24	<i>Pantala flavescens</i>	47	25
<i>Plathemis lydia</i>	40	22	<i>Celithemis elisa</i>	48	30
<i>Ladona julia</i>	40	28	<i>Palpopleura vestita</i>	49	24
<i>Libellula pulchella</i>	41	23	<i>Melammarptis minckii</i>	49	31
<i>Orthemis ferruginea</i>	41	27	<i>Leucorhinia glacialis</i>	51	23
<i>Nannothemis bella</i>	42	32	<i>Macrothemis</i> sp. †.....	52	30
<i>Pseudophlebia minima</i>	43	28			

Suborder **Zygoptera.**

Fam. CALOPTERYGIDÆ.

EPIOPHLEBINÆ. †

<i>Epipophlebia superstes</i>	43	45
-------------------------------------	----	----

EPALLAGINÆ.

<i>Euphæa decorata</i>	51	45*	<i>Anisopleura lestoides</i>	64	54
<i>Rhinocypha</i>	63	61†	<i>Bayadera indica</i>	67	57
<i>Diphlebia lestoides</i>	64	53			

* Minima : second column.

† Maxima : second column.

‡ An undetermined species from Brazil, selected for the extreme reduction of its venation.

§ The preoccupied Selysian name *Palæophlebia* being now replaced by *Epipophlebia* (Calvert, Ent. News, vol. XIV, p. 208, 1903), the subfamily name is here modified to correspond.

VESTALINÆ.

<i>Neurobasis chinensis</i>	46	39	<i>Calopteryx maculata</i>	54	43
<i>Mnais strigata</i>	50	40	<i>Heterina americana</i>	58	40
<i>Calopteryx angustipennis</i>	51	35*	<i>Lais pudica</i>	59	49†

THORINÆ.

<i>Thore picta</i>	52	48	<i>Cora inca</i>	59	50
--------------------------	----	----	------------------------	----	----

Fam. AGRIONIDÆ.

LESTINÆ.

<i>Archilestes grandis</i>	59	38	<i>Lestes rectangularis</i>	66	48
----------------------------------	----	----	-----------------------------------	----	----

AGRIONINÆ.

<i>Calicnemis atkinsoni</i>	57	51	<i>Disparoneura vittata</i>	64	55
<i>Hyponere lugens</i>	58	53	<i>Oxyagrion terminale</i>	64	58
<i>Platycnemis pennipes</i>	61	42	<i>Nehalennia irene</i>	64	61
<i>Hemicnemis bilineata</i>	61	51	<i>Leptagrion macrurum</i>	65	47
<i>Amphiagrion saucium</i>	62	42*	<i>Enallagma doubledayi</i>	65	48
<i>Xanthagrion erythroneurum</i>	62	47	<i>Protoneura capillaris</i>	65	53
<i>Ischnura grællsii</i>	62	48	<i>Telebasis allaudi</i>	65	53
<i>Argia violacea</i>	63	59	<i>Ceriagrion glabrum</i>	65	53
<i>Ceratura capreola</i>	64	48	<i>Acanthagrion gracile</i>	65	54
<i>Mecistogaster</i> sp?.....	64	49	<i>Acanthagrion cheliferrum</i>	65	57
<i>Anomalagrion hastatum</i>	64	51	<i>Erythragrion salvum</i>	67	45
<i>Agriocnemis pulverulans</i>	64	55	<i>Aciagrion pallidum</i>	72	62*

The maxima, minima and averages for both angles are given for the subfamilies, of which representatives were studied, in the following table :

TABLE I.

		Angle of humeral suture.			Angle of tilt of wing bases.					
		Mi.	Ma.	Av.	Mi.	Ma.	Av.			
Anisoptera	{	ÆSCHNIDÆ		ÆSCHININÆ.....	21	42	35	19	34	25
				GOMPHINÆ	31	50	40	22	29	26
				PETALURINÆ	40	—	—	35	—	—
				CORDULEGASTERINÆ.....	37	—	—	35	—	—
	{	LIBELLULIDÆ		MACROMIINÆ.....	29	39	33	18	33	24
				CORDULINÆ.....	30	38	35	23	29	25
				LIBELLULINÆ	28	52	41	19	38	30

* Minima : second column.

† Maxima : second column.

		Angle of humeral suture.			Angle of tilt of wing bases.		
		Mi.	Ma.	Av.	Mi.	Ma.	Av.
Zygoptera	CALOPTERYGIDÆ	EPIOPHLEBIINÆ.....	43	—	—	45	—
		EPALLAGINÆ.....	51	67	62	45	61
		VESTALINÆ.....	46	59	54	35	49
		THORINÆ.....	52	59	55	48	50
	AGRIONIDÆ	LESTINÆ.....	59	66	62	28	48
		AGRIONINÆ.....	54	72	62	42	62
						49	

The above listed material was the best selection that could be made from the material that happened to be at hand in the collection of the senior author. Only papered specimens could be used readily. Whole subfamilies are unrepresented, and the material used is insufficient to furnish a basis for true averages for any of the subfamilies. Yet notwithstanding this, and with all due allowance for error in the making of difficult measurements of angles, some general results are sufficiently evidenced by the figures obtained.

In the first place it is evident that the skewness of the thorax is much greater in the Zygoptera than in the Anisoptera, the average of the former being above the maximum of the latter. The minimum for both angles measured is found in the *Æschninæ*, and the maximum in the *Agrioninæ*. The widest range is shown by the *Libellulinæ*, and it is probable that this is due only in part to the selection of a wider range of representatives of this subfamily.

There was found less correlation between the two angles measured than might have been expected. It will be observed throughout that the maxima and minima rarely fall in the same places in the two columns. The angle that measures the inclination of the humeral suture is with a single exception greater than that measuring the tilt of the wing bases, *Epiophlebia* alone furnishing the reverse condition.

There is little correlation between the size of the insect and the inclination of the humeral suture: just how much, will be seen by glancing down the first column, which in each subfamily is arranged in accordance with the increasing angulation here. There appears to be much more correlation between mere size and the tilt of the wing bases: for it will be noticed in each group that the minimum falls upon one of the largest and the maximum on one of the smallest species.

It must not be forgotten for a moment that these measurements take no account of possible parallelisms within the subfamilies, nor of di-

vergent lines of development, but give merely the degree of progress in two particular lines of specialization arbitrarily selected. The figures, however, are interesting to one who knows these species by sight—even surprising—especially in the Libellulinæ, where they seem to support no one's theory of the interrelationships of the genera.

PHRYGANIDIA CALIFORNICA; *PACKARD*.



BY BEVERLY LETCHER.

Had the general excellence of the account of the life history of this moth as set forth by Vernon L. Kellogg and F. J. Jack (Proceedings of the California Academy of Sciences, second series, Vol. V, 1895, page 562) been maintained throughout their article, there would hardly have been occasion for the present but with them critical observation seems to have ceased with the pupal stage. Other differences are slight but as they are of some interest, may be noted.

To afford a ready comparison of the head measurements, they have been tabulated, those of Kellogg and Jack appearing in the first column and those of the writer in the third. Little value can be attached to the several durations of stages in the fourth column as they were made on larvæ subject to the artificial surroundings of the study: they are from observations of the summer brood while those of Messrs. Kellogg and Jack were made on wintering larvæ.

Stage.	Kellogg and Jack.		Letcher.	
	Mm.	Days.	Mm.	Days.
Egg		10		10
1st	.68	14	.53, .67	8
“ supplementary			.73	8
2d	1.14	13	1.15	6, 8
3d	1.45	17	1.32, 1.47	7
4th	1.88	25	1.82	8
5th	2.21	21	2.20	5
6th	2.57	12	2.31	9
Pupa		10		9

A practical agreement is to be noted for the 1st, 2d, 3d, 4th and 5th stages. My observations show a stage supplemental to the 1st and



Skewness of Thorax in Odonata.